

# Outcomes of Robotic Surgery for Low-Volume Surgeons

 Sridhar Panaiyadiyan,  Rajeev Kumar 

Department of Urology, All India Institute of Medical Sciences, New Delhi, India

## Abstract

When the outcomes are equivalent to the open technique, conventional laparoscopy is a preferred surgical approach because of its minimal invasiveness. However, outcomes following laparoscopy depend on the surgeon's expertise, and there is a significant learning curve to attain efficiency in complex reconstructing laparoscopic procedures. Robotic assistance bridges the gap between open and laparoscopic procedures and allows surgeons with limited laparoscopy experience to offer the benefits of minimally invasive surgery to their patients. While existing data do not show better outcomes with robot assistance compared with laparoscopy for most procedures, these studies are based on data from high-volume surgeons and centers. In reality, a significant number of surgeries are performed by low-volume centers and surgeons, and robotic assistance may enable them to offer benefits of minimally invasive surgery equivalent to those of higher volume centers since robotic assistance is associated with a shorter learning curve than laparoscopy. We review the data on the outcomes of robotic surgery for low-volume surgeons with a focus on centers and surgeons in Asia.

## Introduction

Urology as a specialty has always been at the forefront of technological advancement and has rapidly embraced innovations in devices and techniques. Acceptance of new technology usually depends on its being better for the patient or the health care system[1]. However, being "better" is often associated with being more expensive and thus new technologies must prove themselves to not only give better outcomes but also to do so at a cost that is acceptable for the improvements in outcome at a population level. Being "better for the surgeon" is rarely considered a valid reason for accepting technology that may increase overall costs of treatment.

Among surgical approaches, laparoscopy is universally accepted as the preferred option wherever it has been shown to have outcomes equivalent to open surgery. This is because it provides better cosmesis, lower postoperative pain and earlier recovery; all outcomes that are better for the patient and the health care systems[2]. However, outcomes vary among surgeons and, depending on surgeon expertise, the possible procedures range from simple ablative to complex reconstructive ones that require intracorporeal suturing[3]. Surgical outcomes improve with increasing case load and plateau after a certain level[4]. Once expertise is achieved, outcomes are equivalent irrespective of the surgical approach. However, despite having been available over 30 years, laparoscopy remains challenging for complex reconstructive procedures and has a steep learning curve for such procedures[5].

The introduction of robotic assistance into the surgical armamentarium aimed to bridge the gap between open approaches and laparoscopy and to enable minimally invasive surgery[6]. However, despite having been available now for over 20 years, robotic surgery continues to be questioned and is often derided as a gimmick or marketing tool, primarily because of its high costs, longer operating times, and poor residency training[7,8]. These concerns

### Key Words

Ergonomics, learning curve, robotic assistance, surgical outcomes, surgeon volume, urologic surgery

### Competing Interests

None declared.

### Article Information

Received on April 5, 2022  
Accepted on May 22, 2022  
This article has been peer reviewed.  
Soc Int Urol J. 2022;3(5):323–330  
DOI: 10.48083/PPSC8658

## Abbreviations

EBL	estimated blood loss
LYN	lymph node yield
LPN	laparoscopic partial nephrectomy
OT	operative time
PSM	positive surgical margin
RALP	robotic-assisted laparoscopic pyeloplasty
RAPN	robotic-assisted partial nephrectomy
RARC	robotic-assisted radical cystectomy
RARP	robotic-assisted radical prostatectomy
WIT	warm ischemia time

are fuelled by a lack of studies showing the superiority of robotic assistance over laparoscopy[9]. In the absence of a difference, it is difficult to justify the use of expensive robotic assistance over pure laparoscopy.

Most of this data comes from surgeons and centers that have high volumes. It is only to be expected that once expertise has been acquired through high volumes the outcomes are likely to be similar with each technique. In the real world, most urologists do not have the high volumes of these surgeons or centers[10,11]. However, the patients they treat deserve the same outcomes as those at high-volume centers and it is pertinent to assess whether robotic assistance helps low-volume surgeons provide the benefits of minimally invasive surgery to their patients, which they may not have been able to do through pure laparoscopy.

The learning curve for robot-assisted surgery is shorter than that for conventional laparoscopy in maiden users[12]. Robot assistance allows surgeons trained in open surgery to transition easily to minimally invasive surgery without significant laparoscopy experience, and laparoscopy-naïve surgeons have rapidly gained competence in robot-assisted radical prostatectomy (RARP)[13]. As a corollary, it is plausible that robotic assistance may aid surgeons in a low-volume setting to deliver better outcomes than would be associated with laparoscopy, while providing all the advantages of minimally invasive surgery[11].

With this hypothesis, we reviewed the literature for data on outcomes of robot-assisted surgeries for low-volume surgeons for common urological surgeries such as robot-assisted radical prostatectomy (RARP), robot-assisted partial nephrectomy (RAPN), robot-assisted radical cystectomy (RARC), and robot-assisted laparoscopic pyeloplasty (RALP). This is particularly relevant for Asia since the penetration of robot-assisted surgeries in Asia trails the western world. By 2017, there were 4271 da Vinci Surgical Systems (Intuitive Surgical Inc.) installations worldwide of which 82% were in the United States

Europe as compared with 13% in Asia but with a trend towards increasing Asian installations by 2018[14]. As per the recent data, over 5500 da Vinci Surgical Systems have been installed in 67 countries worldwide[15].

## Low-Volume Robotic Surgeons and/or Hospital

What constitutes “low-volume” for surgeons and hospitals is not clearly defined. Considering RARP as the most common robotic urology surgery, in 2011, Gershman et al. found that 70% of hospitals in the United States were performing less than 50 RARP per year and the complications, hospitalizations, and transfusion rates all declined with increasing experience until around 100 RARPs were being performed per year[16]. While evaluating the impact of surgeon experience, Bravi et al. reported a decline in positive margins from 16.7% to 9.6% with increase of surgeon experience from 10 to 250 procedures[17]. Based on the evaluation of 9810 RARPs in Sweden, Godtman et al.[18] defined very low-volume (surgeon < 13 or hospital < 50 cases), low-volume (surgeon 13 to 25 or hospital 50 to 100 cases), intermediate (surgeon 25 to 50 or hospital 100 to 150 cases), high-volume (surgeon 50 to 76 or hospital 150 to 200 cases) and very high volume (surgeon ≥ 75 or hospital ≥ 200 cases) surgeons and hospitals. Based on the Swedish national guidelines for prostate cancer, Godtman et al. defined these cut-off values as a multiple of the minimum number of RARPs that should be performed by the surgeon and the center. It is expected that these numbers would differ for other procedures. This lack of consensus on the definition of low-volume surgeon would be related to the different learning curve in achieving a benchmark optimum outcome.

## Methods

We searched PubMed/MEDLINE, Embase, Web of Science, and Google Scholar databases for relevant studies published in English using the following MeSH key words: (robotic urology surgery OR robotic radical prostatectomy OR RARP OR radical cystectomy OR RARC OR partial nephrectomy OR pyeloplasty) AND (low-volume surgeons OR low-volume center OR learning curve). The abstracts and full articles of systematic reviews and meta-analyses were reviewed. Case reports, letters, short communications, and articles in languages other than English language were excluded.

## Radical Prostatectomy

Reviewing the outcomes of radical prostatectomies done at high-volume centers, Coelho et al. showed that RARP delivered better outcomes in terms of weighted mean PSM, continence, and potency rates compared with retropubic and laparoscopic radical prostatectomies[19]. However, with the increasing adoption of robotic system in the recent years, there has been an increase in the

number of RARP being done at low-volume centers[20]. It is now estimated that between 33% and 70% of RARP in the United States are done by low-volume surgeons[10].

With conventional laparoscopic radical prostatectomy, the number needed to attain a 90% recurrence-free probability is estimated to be 750 cases[21]. In contrast, with RARP, about 80 to 120 cases are needed to attain a comparable perioperative, oncological, and functional outcomes[22]. Further, with increasing experience in RARP, the perioperative complications declined from 9.8% in low-volume surgeons to 6.7% in intermediate-volume surgeons and to 2.2% in high-volume surgeons[23]. In a retrospective analysis from India, Garg et al. showed an inverse stage migration pattern in their RARP practice. Over the time, with increasing experience in RARP, patients with more unfavorable disease characteristics underwent RARP[24].

A recent multi-national and multicenter retrospective study discussed the RARP outcomes (n = 207) of 3 fellowship-trained surgeons from low-volume regions of Gulf Cooperation Council countries[25]. This study showed trifecta outcomes comparable to those of international centers. At the 12-month follow-up, 35.8% of patients were potent, 94.6% were continent, and 9.2% had biochemical recurrence[25]. In a recent study from low-intermediate volume center, Afferi et al. retrospectively analyzed 604 men who underwent nerve-sparing RARP[26]. At a median follow-up of 28 months, the authors reported an increase in the pentapecta outcomes from 38% to 44%, with erectile dysfunction being the main limiting factor for non-achievement of pentapecta in 71% cases. However, in high-risk prostate cancers, biochemical recurrence limited the pentapecta achievement (61%)[26].

## Partial Nephrectomy

The current standard of care for small renal masses is partial nephrectomy[27]. Prior reports showed equivalent oncological outcomes between open and laparoscopic approaches for partial nephrectomies[28,29]. However, laparoscopic partial nephrectomy (LPN) is a technically demanding procedure requiring intracorporeal suturing with simultaneous consideration of warm ischemia time (WIT) to prevent ischemic renal injury[30]. Considering that laparoscopic radical nephrectomy was easier to perform, it was hypothesized that partial nephrectomy was underutilized for eligible candidates[31]. After successful incorporation of robotic assistance into radical prostatectomy, robot-assisted partial nephrectomy (RAPN) was quickly and widely adopted[32]. A surgeon needs to perform fewer RAPN than LPN to attain a comparable level of

competency[33]. It is estimated that about 565 cases are required to achieve the target WIT (< 20 to 25 minutes) in LPN[34], whereas it requires about 20 and 50 cases, for RAPN for equivalent WIT and operative time (OT), respectively[5]. In contrast, expert renal surgeons require < 30 cases of RAPN to achieve an equivalent proficiency.

In a low-volume center in Kuwait, a fellowship-trained robotic surgeon retrospectively assessed the perioperative, trifecta, and pentapecta outcomes following 43 RAPN cases performed over 6 years. Operative time and estimated blood loss (EBL) improved significantly after the first quartile of patients (n = 14). Trifecta and pentapecta outcomes were achieved in 93% and 81.8% cases, respectively. A higher number of complex cases with RENAL nephrometry scores of 7 to 12 were performed in the second (n = 14) and third (n = 15) quartiles[35].

In a prospective study, Dias et al. reported outcomes of RAPN by a laparoscopy trained surgeon with no prior robotic surgery experience. In 108 cases performed over 5 years, WIT ≤ 20 minutes, OT ≤ 150 minutes, and EBL < 100mL were achieved after 44, 44, and 54 cases, respectively. A trifecta outcome was achieved in 67.6% cases[36]. Motoyama et al. in a retrospective study reported the outcomes of 65 RAPN performed by a surgeon with no prior LPN experience but with extensive robotic surgery experience[37]. The learning curve was analyzed by dividing cases into 5 groups of 13 consecutive patients. Trifecta outcomes were achieved in 89.2% cases. The authors reported a WIT of ≤ 20 minutes and console time of ≤ 150 minutes could be achieved at fourth and sixth procedures, respectively[37].

In terms of complications, a multicenter study showed that a low-volume surgeon performing < 7 cases of RAPN per year had a complication rate (18.1% versus 15.9% versus 16.1%, *P* = 0.81) comparable to high-volume (15 to 30 RAPN cases/year) and very high-volume (> 30 RAPN cases/year) surgeons[38]. In a recent database study involving 124 institutions in Japan, Yokoyama et al. reported the trends and safety of RAPN done at low-volume (n = 616) and high-volume (n = 3106) institutions. After propensity score matching, the authors did not find any significant differences in anesthesia time, blood transfusions, and complication rates between low-volume and high-volume centers[39]. Similarly, data from a single-center study in Turkey confirmed that a comparable trifecta outcome (77%) can be achieved by low-volume surgeon[40].

## Radical Cystectomy

Radical cystectomy with urinary diversion has one of the highest morbidity rates among urological procedures, with postoperative complication rates of up to 69%[41]. Minimally invasive surgery, both pure laparoscopy

and robot-assisted, are feasible alternatives to open radical cystectomy. Data from the RAZOR trial suggest that RARC is associated with decreased blood loss, transfusion rate, and hospital stay compared with the open approach[41]. Further, RARC has been shown to be non-inferior to the open approach in terms of 3-year progression-free and overall survival[42]. The learning curve for such complex procedures is estimated to be 21 to 30 cases when using benchmarks of operating time of < 6.5 hours, lymph node yield (LNY) of 20, and < 5% positive surgical margin (PSM). The number needed to reach the target OT was 21 cases. A total of 8, 20, and 30 cases were required to attain a LNY of 12, 16, and 20, respectively. Similarly, 30 cases were required to reach competency for PSM[43].

Guru et al. divided 100 consecutive RARC procedures into 4 quartiles. The overall mean OT and EBL were 343 minutes and 598 mL, respectively. The mean OT reduced from 375 minutes in first quartile to 352 minutes in the last quartile. Similarly, the LNY increased from 14 nodes in first quartile to 23 nodes in the last quartile. While there were 4 cases of PSM in the first quartile, none of the patients in the last quartile had a positive margin. The authors reported no difference in the complication rate over time[44].

Experience in one robot-assisted procedure helps improve outcomes in other procedures. Hayn et al., using the International Robotic Cystectomy Consortium database, report that surgeons with prior experience in RARP provide better RARC outcomes with significant differences in OT, EBL, and LNY between surgeons with < 50 RARPs and those with 51 to 100 RARPs[45].

The adoption of RARC has been increasing in Asian countries in recent years. A single-center retrospective analysis from India of the initial experience of 63 RARC with extracorporeal urinary diversion reported a mean OT, EBL, and hospital stay of 348 minutes, 868 mL, and 10.4 days, respectively. The mean LNY was 12.4, ranging between 3 and 25 nodes[46]. These data are very similar to those reported from larger centers[44].

## Pyeloplasty

Robotic assistance has also helped lower the steep learning curve associated with laparoscopic pyeloplasty for uretero-pelvic junction obstruction[47]. It has been documented to be of particular advantage in patients with anatomical variations like lower moiety uretero-pelvic junction obstruction, malrotated kidneys, and secondary uretero-pelvic junction obstruction[48–52].

RALP is a relatively simple procedure and has been considered as a gateway to advanced reconstructive procedures[53]. A novice can achieve an operating time within 1 standard deviation of the open pyeloplasty

**TABLE 1.**

Learning curve of common robotic urological procedures

Procedure*	Number of cases required	Outcome measures
RARPa	80–120 cases[21]	Perioperative, oncological, and functional outcomes
RAPN	20 cases[5] 50 cases[5]	WIT OT
RARC	21–30 cases[41]	OT < 6.5 hours; LNY of 20; PSM < 5%
RALP	15–20 cases[52]	OT

*RARP: robotic-assisted radical prostatectomy; RAPN: robotic-assisted partial nephrectomy; RARC: robotic-assisted radical cystectomy; RALP: robotic-assisted laparoscopic pyeloplasty; WIT: warm ischemia time; OT: operative time; LNY: lymph node yield and PSM: positive surgical margin*

*\*All the data provided here are originated from surgeons trained in both open and laparoscopy except (a) Denotes the learning curve of a surgeon with no prior laparoscopic experience.*

after 15 to 20 cases[54]. However, it takes only about 5 cases for a surgeon expert in open procedures[55]. It is not surprising that robotic assistance does not offer any benefit to experienced laparoscopic surgeons with good intracorporeal suturing skills[56]. However, a relatively faster learning curve could prompt low-volume surgeons to offer RALP to their patients to deliver comparable outcomes.

## Ergonomics

The ergonomics of the procedures for surgeons are often under-reported, and poor ergonomics affect physical, mental, and economic well-being. A greater degree of adverse ergonomic problems are expected among urologists who perform a wide range of operative procedures such as open, laparoscopic, robotic, endoscopic, and microscopic surgeries[57]. Catanzarite et al. have shown that work-related musculoskeletal disorders are prevalent in 66% to 90%, 73% to 100%, and 23% to 80% for open, laparoscopic, and robotic surgeries, respectively[58]. Surgeon fatigue after performing a conventional laparoscopic pyeloplasty may limit the number of procedures that may be performed on the same day[59]. Specifically, post-procedure fatigue affects precise intracorporeal suture placement[60]. Although

robot-assisted surgery appears to be ergonomically better, it too can lead to a strain on trunk, wrist, and fingers[58]. For instance, one of the most common ergonomic issues while performing robotic surgery is abducted shoulder and not maintaining a right angle with the biceps, resulting in lifting off the elbows from the armrest. Surgeons have to use the clutch control to mitigate the abnormal posture[57].

The learning curves of common robotic urological procedures are shown in **Table 1**. These numbers should be interpreted with some caution as the data could be heterogenous as the data include different cohorts. Further, various confounders affecting the outcomes of different robotic procedures were not addressed. However, our work provides an overview of data on outcomes of common robot-assisted surgeries for low-volume surgeons.

### Role of Various Robotic Training Models

Various training models have the potential to help low-volume robotic surgeons deliver a reasonable surgical outcome without compromising patient safety. In this context, expert proctorship helped trainees achieve efficiency and early performance of independent robotic procedures. With the introduction of dual-console da Vinci Si or Xi Surgical systems, the proctorship program has become more convenient for both proctor and trainee[61–63]. As the technology advances, the future

looks promising for telementoring, which may facilitate remote proctorship[64]. A fellowship program has been considered the most structured robotic training model as it involves a systematic method of training[65]. The trainees perform progressive steps of the operation in succession under the guidance of an expert robotic surgeon with constant and immediate feedback[61]. Lastly, simulation-based training allows trainees to develop surgical skills without risking patient safety. It provides an objective assessment of particular surgical skills. In recent years, various robotic simulation platforms have been developed and are commercially available. In particular, the da Vinci skills simulator is closely attached to the surgeon console that would be otherwise used for robotic-assisted surgeries[66].

### Conclusion

Robotic assistance has the potential to help low-volume surgeons and hospitals offer minimally invasive surgery to their patients. Robotic assistance enables surgeons to reach a higher degree of competence more quickly and easily. While the costs associated with robot assistance continue to be high, the potential benefit of minimally invasive surgery over open surgery would help mitigate concerns. Ergonomic benefits for the surgeon are unmeasured, but may also be a significant advantage of robotic-assisted surgery.

### References

- Venkatramani V. Urovision 2020: The future of urology. *Indian J Urol*.2015;31(2):150-155. DOI: 10.4103/0970-1591.152918
- Buia A, Stockhausen F, Hanisch E. Laparoscopic surgery: a qualified systematic review. *World J Methodol*.2015;5(4):238-254. DOI: 10.5662/wjm.v5.i4.238
- Kumar R, Hemal AK. Emerging role of robotics in urology. *J Minim Access Surg*.2005;1(4):202-210. DOI: 10.4103/0972-9941.19268
- Vincent C, Moorthy K, Sarker SK, Chang A, Darzi AW. Systems approaches to surgical quality and safety: from concept to measurement. *Ann Surg*.2004;239(4):475-482. DOI: 10.1097/01.sla.0000118753.22830.41
- Abboudi H, Khan MS, Guru KA, Froghi S, de Win G, Van Poppel H, et al. Learning curves for urological procedures: a systematic review: Learning curves for urological procedures. *BJU Int*.2014;114(4):617-629. DOI: 10.1111/bju.12315
- Shah J, Vyas A, Vyas D. The history of robotics in surgical specialties. *Am J Robot Surg*.2014;1(1):12-20. DOI: 10.1166/ajrs.2014.1006
- Ashrafian H, Clancy O, Grover V, Darzi A. The evolution of robotic surgery: surgical and anaesthetic aspects. *Br J Anaesth*.2017;119(suppl\_1):i72-84. DOI: 10.1093/bja/aex383
- Farivar BS, Flannagan M, Leitman IM. General surgery residents' perception of robot-assisted procedures during surgical training. *J Surg Educ*.2015;72(2):235-42. DOI: 10.1016/j.jsurg.2014.09.008
- Dhanani NH, Olavarria OA, Bernardi K, Lyons NB, Holihan JL, Loor M, et al. The evidence behind robot-assisted abdominopelvic surgery: a systematic review. *Ann Intern Med*.2021;174(8):1110-7. DOI: 10.7326/M20-7006
- Wilt TJ, Shamlilian TA, Taylor BC, MacDonald R, Kane RL. Association between hospital and surgeon radical prostatectomy volume and patient outcomes: a systematic review. *J Urol*.2008;180(3):820-828; discussion 828-829. DOI: 10.1016/j.juro.2008.05.010
- Panaiyadiyan S, Kumar R. Robotic assistance helps low-volume surgeons deliver better outcomes to their patients. *Indian J Urol*.2021;37(1):1. DOI: 10.4103/iju.IJU\_216\_20
- Anderberg M, Larsson J, Kockum CC, Arnbjörnsson E. Robotics versus laparoscopy - an experimental study of the transfer effect in maiden users. *Ann Surg Innov Res*.2010;4(1):3. DOI: 10.1186/1750-1164-4-3
- Ahlering TE, Skarecky D, Lee D, Clayman RV. Successful transfer of open surgical skills to a laparoscopic environment using a robotic interface: initial experience with laparoscopic radical prostatectomy. *J Urol*.2003;170(5):1738-1741. DOI: 10.1097/01.ju.0000092881.24608.5e

14. Azhar RA, Elkoushy MA, Aldousari S. Robot-assisted urological surgery in the Middle East: Where are we and how far can we go? *Arab J Urol*.2019;17(2):106-113. DOI: 10.1080/2090598X.2019.1601003
15. Koukourikis P, Rha KH. Robotic surgical systems in urology: What is currently available? *Investig Clin Urol*.2021;62(1):14-22. DOI: 10.4111/icu.20200387
16. Gershman B, Meier SK, Jeffery MM, Moreira DM, Tollefson MK, Kim SP, et al. Redefining and contextualizing the hospital volume-outcome relationship for robot-assisted radical prostatectomy: implications for centralization of care. *J Urol*.2017;198(1):92-99. DOI: 10.1016/j.juro.2017.01.067
17. Bravi CA, Tin A, Vertosick E, Mazzone E, Martini A, Dell'Oglio P, et al. The impact of experience on the risk of surgical margins and biochemical recurrence after robot-assisted radical prostatectomy: a learning curve study. *J Urol*.2019;202(1):108-113. DOI: 10.1097/JU.000000000000147
18. Godtman RA, Persson E, Cazzaniga W, Sandin F, Carlsson S, Ahlgren G, et al. Association of surgeon and hospital volume with short-term outcomes after robot-assisted radical prostatectomy: Nationwide, population-based study. *PLoS One*.2021;16(6):e0253081. DOI: 10.1371/journal.pone.0253081
19. Coelho RF, Rocco B, Patel MB, Orvieto MA, Chauhan S, Ficarra V, et al. Retropubic, laparoscopic, and robot-assisted radical prostatectomy: a critical review of outcomes reported by high-volume centers. *J Endourol*.2010;24(12):2003-2015. DOI: 10.1089/end.2010.0295
20. Leow JJ, Leong EK, Serrell EC, Chang SL, Gruen RL, Png KS, et al. Systematic review of the volume-outcome relationship for radical prostatectomy. *Eur Urol Focus*.2018;4(6):775-789. DOI: 10.1016/j.euf.2017.03.008
21. Vickers AJ, Savage CJ, Hruza M, Tuerk I, Koenig P, Martínez-Piñero L, et al. The surgical learning curve for laparoscopic radical prostatectomy: a retrospective cohort study. *Lancet Oncol*. 2009;10(5):475-480. DOI: 10.1016/S1470-2045(09)70079-8
22. Gumus E, Boylu U, Turan T, Onol FF. The learning curve of robot-assisted radical prostatectomy. *J Endourol*.2011;25(10):1633-1637. DOI: 10.1089/end.2011.0071
23. Hirasawa Y, Yoshioka K, Nasu Y, Yamamoto M, Hinotsu S, Takenaka A, et al. Impact of surgeon and hospital volume on the safety of robot-assisted radical prostatectomy: a multi-institutional study based on a national database. *Urol Int*.2017;98(3):334-342. DOI: 10.1159/000460304
24. Garg H, Seth A, Singh P, Kumar R. Changing trends in robot-assisted radical prostatectomy: Inverse stage migration-a retrospective analysis. *Prostate Int*.2021;9(3):157-162. DOI: 10.1016/j.pnrl.2021.04.002
25. Azhar RA, Aldousari S, Alghamdi MM, Alotaibi MF, Alkhateeb SS, Nassir AM, et al. Robot-assisted radical prostatectomy in low-volume regions: should it be abandoned or adopted? A multi-institutional outcome study. *J Endourol*.2021;35(7):1013-1019. DOI: 10.1089/end.2020.0770
26. Afferi L, Moschini M, Baumeister P, Zamboni S, Cornelius J, Ineichen G, et al. Trends in risk-group distribution and Pentafecta outcomes in patients treated with nerve-sparing, robot-assisted radical prostatectomy: a 10-year low-intermediate volume single-center experience. *World J Urol*. 2021;39(2):389-397. DOI: 10.1007/s00345-020-03206-y
27. Miller DC, Hollingsworth JM, Hafez KS, Daignault S, Hollenbeck BK. Partial nephrectomy for small renal masses: an emerging quality of care concern? *J Urol*.2006;175(3 Pt 1):853-857; discussion 858. DOI: 10.1016/S0022-5347(05)00422-2
28. You C, Du Y, Wang H, Peng L, Wei T, Zhang X, et al. Laparoscopic versus open partial nephrectomy: a systemic review and meta-analysis of surgical, oncological, and functional outcomes. *Front Oncol*.2020;10:583979. DOI: 10.3389/fonc.2020.583979
29. Yu Y, Wang W, Xiong Z, Yang Z, Li J, Shen Y, et al. Comparison of perioperative outcomes between laparoscopic and open partial nephrectomy for different complexity renal cell carcinoma based on the R.E.N.A.L. nephrometry score. *CMAR*.2021;13:7455-7461. DOI: 10.2147/CMAR.S324457
30. Jain S, Gautam G. Robotics in urologic oncology. *J Minim Access Surg*.2015;11(1):40-44. DOI: 10.4103/0972-9941.147687
31. Abouassaly R, Alibhai SMH, Tomlinson G, Timilshina N, Finelli A. Unintended consequences of laparoscopic surgery on partial nephrectomy for kidney cancer. *J Urol*.2010;183(2):467-472. DOI: 10.1016/j.juro.2009.10.002
32. Cheung H, Wang Y, Chang SL, Khandwala Y, Del Giudice F, Chung BI. Adoption of robot-assisted partial nephrectomies: a population-based analysis of U.S. surgeons from 2004 to 2013. *J Endourol*.2017;31(9):886-892. DOI: 10.1089/end.2017.0174
33. Hanzly M, Frederick A, Creighton T, Atwood K, Mehedint D, Kauffman EC, et al. Learning curves for robot-assisted and laparoscopic partial nephrectomy. *J Endourol*.2015;29(3):297-303. DOI: 10.1089/end.2014.0303
34. Gill IS, Kamoi K, Aron M, Desai MM. 800 Laparoscopic partial nephrectomies: a single surgeon series. *J Urol*.2010;183(1):34-41. DOI: 10.1016/j.juro.2009.08.114
35. Aldousari S, Yaiesh S, Alkandari O. Total robotic surgical volume influences outcomes of low-volume robotic-assisted partial nephrectomy over an extended duration. *CUAJ*.2021;15(9). DOI: 10.5489/cuaj.6880
36. Dias B, Ali M, Dubey S, Krishnaswamy S, Rao A, Dubey D. Impact of learning curve on the perioperative outcomes following robot-assisted partial nephrectomy for renal tumors. *Indian J Urol*.2018;34(1):62. DOI: 10.4103/iju.IJU\_169\_17
37. Motoyama D, Matsushita Y, Watanabe H, Tamura K, Suzuki T, Ito T, et al. Initial learning curve for robot-assisted partial nephrectomy performed by a single experienced robotic surgeon. *Asian J Endosc Surg*.2020;13(1):59-64. DOI: 10.1111/ases.12683

38. Peyronnet B, Tondut L, Bernhard J-C, Vaessen C, Doumerc N, Sebe P, et al. Impact of hospital volume and surgeon volume on robot-assisted partial nephrectomy outcomes: a multicentre study. *BJU Int*.2018;121(6):916-922. DOI: 10.1111/bju.14175
39. Yokoyama M, Ishioka J, Toba M, Fukushima H, Tanaka H, Yoshida S, et al. Trends and safety of robot-assisted partial nephrectomy during the initial 2-year period after government approval in Japan: a nationwide database study from 2016 to 2018. *Int J Urol*.2021;28(12):1268-1272. DOI: 10.1111/iju.14698
40. Basatac C, Akpınar H. « Trifecta » outcomes of robot-assisted partial nephrectomy: Results of the « low volume » surgeon. *Int Braz J Urol*.2020;46(6):943-949. DOI: 10.1590/S1677-5538.IBJU.2019.0396
41. Parekh DJ, Reis IM, Castle EP, Gonzalgo ML, Woods ME, Svatek RS, et al. Robot-assisted radical cystectomy versus open radical cystectomy in patients with bladder cancer (RAZOR): an open-label, randomised, phase 3, non-inferiority trial. *Lancet*.2018;391(10139):2525-2536. DOI: 10.1016/S0140-6736(18)30996-6
42. Venkatramani V, Reis IM, Castle EP, Gonzalgo ML, Woods ME, Svatek RS, et al. Predictors of recurrence, and progression-free and overall survival following open versus robotic radical cystectomy: analysis from the RAZOR Trial with a 3-Year followup. *J Urol*.2020;203(3):522-529. DOI: 10.1097/JU.0000000000000565
43. Hayn MH, Hussain A, Mansour AM, Andrews PE, Carpentier P, Castle E, et al. The Learning curve of robot-assisted radical cystectomy: results from the International Robotic Cystectomy Consortium. *Eur Urol*.2010;58(2):197-202. DOI: 10.1016/j.eururo.2010.04.024
44. Guru KA, Perlmutter AE, Butt ZM, Piacente P, Wilding GE, Tan W, et al. The learning curve for robot-assisted radical cystectomy. *JSLS*.2009;13(4):509-14. DOI: 10.4293/108680809X12589998404128
45. Hayn MH, Hellenthal NJ, Hussain A, Andrews PE, Carpentier P, Castle E, et al. Does previous robot-assisted radical prostatectomy experience affect outcomes at robot-assisted radical cystectomy? Results from the International Robotic Cystectomy Consortium. *Urology*.2010;76(5):1111-1116. DOI: 10.1016/j.urol.2010.05.010
46. Shrivastava N, Nayak B, Dogra P, Kumar R, Singh P. Robot-assisted laparoscopic radical cystectomy with extracorporeal urinary diversion: Initial experience and outcomes. *Indian J Urol*.2018;34(2):122-126. DOI: 10.4103/iju.IJU\_65\_17
47. Dothan D, Raisin G, Jaber J, Kocherov S, Chertin B. Learning curve of robotic-assisted laparoscopic pyeloplasty (RALP) in children: how to reach a level of excellence? *J Robotic Surg*.2021;15(1):93-97. DOI: 10.1007/s11701-020-01082-7
48. Hemal AK, Mishra S, Mukharjee S, Suryavanshi M. Robot assisted laparoscopic pyeloplasty in patients of ureteropelvic junction obstruction with previously failed open surgical repair. *Int J Urol*.2008;15(8):744-746. DOI: 10.1111/j.1442-2042.2008.02091.x
49. Nayyar R, Gupta NP, Hemal AK. Robotic management of complicated ureteropelvic junction obstruction. *World J Urol*.2010;28(5):599-602. DOI: 10.1007/s00345-009-0469-y
50. Bora GS, Parmar K, Mavuduru RS. Robot-assisted pyeloplasty for pelvi-ureteric junction obstruction of lower moiety in partial duplex system: a technical challenge. *Indian J Urol*.2016;32(4):314-316. DOI: 10.4103/0970-1591.191264
51. Numakura K, Muto Y, Saito M, Narita S, Inoue T, Habuchi T. Robot-assisted laparoscopic pyeloplasty for ureteropelvic junction obstruction with duplex system. *Urol Case Rep*.2020;30:101138. DOI: 10.1016/j.eucr.2020.101138
52. Dirie NI, Ahmed MA, Mohamed MA, Zhang Z, Wang S. Robot-assisted laparoscopic pyeloplasty in adults: a comparison analysis of primary versus redo pyeloplasty in a single center. *Urol J*.2020;18(1):45-50. DOI: 10.22037/uj.v16i7.5257
53. Tasian GE, Casale P. The robotic-assisted laparoscopic pyeloplasty: gateway to advanced reconstruction. *Urol Clin North Am*.2015;42(1):89-97. DOI: 10.1016/j.ucl.2014.09.008
54. Sorensen MD, Delostrinos C, Johnson MH, Grady RW, Lendvay TS. Comparison of the learning curve and outcomes of robotic assisted pediatric pyeloplasty. *J Urol*.2011;185(6 Suppl):2517-2522. DOI: 10.1016/j.juro.2011.01.021
55. Bowen DK, Lindgren BW, Cheng EY, Gong EM. Can proctoring affect the learning curve of robotic-assisted laparoscopic pyeloplasty? Experience at a high-volume pediatric robotic surgery center. *J Robotic Surg*.2017;11(1):63-7. DOI: 10.1007/s11701-016-0613-9
56. Maheshwari P, Arora A. Robot-assisted pyeloplasty: The way forward. *J Min Access Surg*. 2021;17(2):274. DOI: 10.4103/jmas.JMAS\_59\_20
57. Gabrielson AT, Clifton MM, Pavlovich CP, Biles MJ, Huang M, Agnew J, et al. Surgical ergonomics for urologists: a practical guide. *Nat Rev Urol*.2021;18(3):160-169. DOI: 10.1038/s41585-020-00414-4
58. Catanzarite T, Tan-Kim J, Whitcomb EL, Menefee S. Ergonomics in surgery: a review. *Female Pelvic Med Reconstr Surg*.2018;24(1):1-12. DOI: 10.1097/SPV.0000000000000456
59. Kumar R, Nayak B. Robotic versus conventional laparoscopic pyeloplasty: a single surgeon concurrent cohort review. *Indian J Urol*.2013;29(1):19-21. DOI: 10.4103/0970-1591.109978
60. Margossian H, Garcia-Ruiz A, Falcone T, Goldberg JM, Attaran M, Miller JH, et al. Robotically assisted laparoscopic tubal anastomosis in a porcine model: a pilot study. *J Laparoendosc Adv Surg Tech A*.1998;8(2):69-73. DOI: 10.1089/lap.1998.8.69
61. Santok GD, Raheem AA, Kim LH, Chang K, Chung BH, Choi YD, et al. Proctorship and mentoring: Its backbone and application in robotic surgery. *Investig Clin Urol*.2016;57(Suppl 2):S114. DOI: 10.4111/icu.2016.57.S2.S114
62. Chowdhury A, Tan LGL, Chiong E, Rha KH, Tiong HY. Transitioning to robotic partial nephrectomy with a team-based proctorship achieves the desired improved outcomes over open and laparoscopic partial nephrectomy. *Updates Surg*.2021;73(3):1189-1196. DOI: 10.1007/s13304-021-01028-0

- 
63. Mason MD, Herndon CDA, Herbst KW, Poston TL, Brandt EJ, Peters CA, et al. Proctor environment facilitates faculty training in pediatric robotic-assisted laparoscopic pyeloplasty. *J Robotic Surg.* 2014;8(4):365-369. DOI: 10.1007/s11701-014-0481-0
64. Santomauro M, Reina GA, Stroup SP, L'Esperance JO. Telementoring in robotic surgery. *Curr Opin Urol.* 2013;23(2):141-5. DOI: 10.1097/MOU.0b013e32835d4cc2
65. El Sherbiny A, Eissa A, Ghaith A, Morini E, Marzotta L, Sighinolfi MC, et al. Training in urological robotic surgery. Future perspectives. *Arch Esp Urol.* 2018;71(1):97-107.
66. Canalichio KL, Berrondo C, Lendvay TS. Simulation training in urology: state of the art and future directions. *Adv Med Educ Pract.* 2020;11:391-396. DOI: 10.2147/AMEP.S198941