

Supplementary Appendix 2. Functional outcomes at 3-, 6-, 12-, and 24-month follow-up visits

Clinical & paraclinical data	Preoperative N = 1174	3 months N = 954	6 months N = 406	12 months N = 408	24 months N = 232
IPSS	19 [13-23]	5 [3-9]	5 [2-9]	3 [1-6]	5 [2-8]
IPSS Q8	5 [4-6]	1 [0-3]	1 [0-2]	1 [0-2]	1 [0-2]
IIEF5	13.7 ± 8.3	13.2 ± 8.3	12 ± 8.2	12.9 ± 8.7	11.2 ± 8.6
Qmax	9.5 ± 4.6	22.9 ± 12.7	23.7 ± 13.2	24.8 ± 12.6	24.7 ± 13.3
PVR	150.5 ± 188.4	19.5 ± 51.6	16.3 ± 33.6	27.7 ± 55.9	41.1 ± 74.2
PSA	6.8 ± 9.4	1.8 ± 7.7	1.8 ± 2.8	1.8 ± 3.3	3.1 ± 6.5
Urinary Incontinence	6 (0.5)	269 (22.9)	75 (6.4)	107 (9.1)	30 (2.5)
Urinary protection	3 (0.2)	136 (11.6)	45 (3.8)	31 (2.6)	25 (2.1)
<i>Values expressed as mean (SD), median [interquartile range] or n (%).</i>					

Klein C, Marquette T, Capon G, Alezra E, Blanc P, Estrade V, et al. Implementing HoLEP in an Academic Department With Multiple Surgeons in Training: Mentoring Is the Key for Success. Soc Int Urol J. 2023;4(1):11–18. doi: 10.48083/UJCR1584