

# Harm Related to Social Media Misinformation on Pelvic Organ Prolapse in YouTube, Instagram, and Pinterest Posts

Chaoyang Wang,<sup>✉1</sup> Juhye Kang,<sup>1</sup> Emily Gerard,<sup>1</sup> Stacy Loeb,<sup>2</sup> Rena D. Malik<sup>1</sup>

<sup>1</sup>Department of Urology at the University of Maryland School of Medicine, Baltimore, United States

<sup>2</sup>Department of Urology at NYU Langone Health, New York City, United States

## Abstract

Social media can improve patient education but may pose risks due to misinformation. There is no consensus on categorizing types of misinformation and harm. This study aimed to categorize and quantify misinformation and resultant harm from posts on YouTube, Pinterest, and Instagram on pelvic organ prolapse (POP), a prevalent benign condition that impacts quality of life. We conducted a descriptive study of 300 posts presented in these platforms in 2019. Using Fisher exact test, we show a significant difference in the distribution of misinformation between social media platforms. Harmful posts were most frequently present on Pinterest, leading to harmful inaction and economic harm.

## Introduction

In the United States, 8 in 10 internet users search for healthcare information online, a majority using social media[1]. A recent study examining misinformation in online cancer resources concluded that 32.5% of these articles contained misinformation and the majority were deemed harmful[2]. While misinformation about malignancies can pose significant health risks, it remains unclear whether misinformation about prevalent benign conditions such as pelvic organ prolapse (POP) leads to harm.

With growing numbers of social media platforms, instruments to categorize diverse social media posts regarding the type of misinformation and potential harm may be challenging to implement due to the vastly different content format and length presented in these platforms. Therefore, a streamlined classification of the type and nature of health-related misinformation compatible across multiple networks is needed.

POP is a benign condition that impacts quality of life with a prevalence as high as 50% of women, with 12% electing for surgical correction[3]. We previously identified the presence of “low to moderate quality” information in 74.1% of posts on pelvic organ prolapse on Pinterest, YouTube, and Instagram[4].

We investigated 3 prominent social media platforms (YouTube, Pinterest, and Instagram) for POP posts to categorize the quantity and types of misinformation represented, as well as the degree and type of potential harm resulting from exposure to misinformation about this benign condition.

### Key Words

Social media, misinformation, harm, pelvic organ prolapse

### Competing Interests

None declared.

### Article Information

Received on, October 4, 2022  
Accepted on, November 26, 2022  
This article has been peer reviewed.  
Soc Int Urol J. 2023;4(2):131–135  
DOI: 10.48083/PGWG4918

## Materials and Methods

We conducted a descriptive study in October 2021 that was exempt by the University of Maryland Internal Review Board (IRB). Relevant YouTube videos and Pinterest and Instagram posts were identified by searching for “pelvic organ prolapse” on YouTube and Pinterest and “#PelvicOrganProlapse” on Instagram in September 2019. These platforms were chosen because they are among the top 15 most used social media platforms and have a proprietary search engine to identify relevant and publicly available pelvic organ prolapse posts. The first 100 posts on each platform were analyzed for quality, understandability, actionability, and misinformation as detailed by our previous study[4]. Misinformation was assessed using a Likert scale from 1 (no misinformation) to 5 (high misinformation).

We reviewed posts with misinformation scores  $\geq 2$  for type and amount of misinformation, and the type of resultant harm[2]. To measure harmfulness, reviewers used a Likert scale from 1 (certainly not harmful) to 5 (certainly harmful). Harm was defined as harmful inaction (encouragement to forgo standard of care), economic harm (money spent on ineffective treatments), harmful action (toxic effects of the suggested test or treatment), and harmful interaction (medical interactions with curative therapies). Type of misinformation was defined based on prior work: fabricated content (a completely false statement), misleading content (misleading use of content to frame an issue), imposter content (genuine sources impersonated with false sources), manipulative content (genuine information or imagery manipulated to deceive), false connection (headlines, visuals, or captions that do not support the content), and false context (genuine content shared with false contextual information)[5]. Each social media post was reviewed by at least 2 reviewers. Disagreements were reviewed until consensus was achieved.

Descriptive statistics were calculated as the number of observations and percentages. Data were analyzed using Fisher exact test and examined for associations between the categorical variables of interest: social media platform, quantity of misinformation, and harmfulness of medical claims. *P*-values  $< 0.05$  were significant.

## Results

Of 300 total posts including 100 posts per platform, 82 posts were identified as misinformative comprising 18 YouTube videos, 24 Pinterest posts, and 40 Instagram posts. Instagram and Pinterest had a greater proportion of posts that contained “mostly false” claims, whereas YouTube had a greater proportion of videos with “mostly true” claims. “Mostly false” and “false” medical claims

regarding POP were identified in 10 of 82 (12.2%) of the posts (0% YouTube videos, 15% Instagram posts, and 17% Pinterest posts).

Social media misinformation about POP most frequently arises from fabricated content (4 of 10), false connection (5 of 10), or misleading content (6 of 10). For example, one video included chemotherapy and radiation as a nonsurgical treatment option for POP (fabricated content) and another post misrepresented an article published in the journal *Nature* to claim that starvation and dysbiosis were prominent risk factors for POP (false connection).

Of 82 misinformative posts, 8.5% were found to be “probably harmful” (6% YouTube videos, 3% Instagram posts, and 21% Pinterest posts). Among these 7 posts, harmful inaction (5 of 7 posts) and economic harm (5 of 7 posts) were the most common types of harm, suggesting that harmful misinformation may result in delays of care and/or more expensive alternative routes of treatment (Table 1). For example, one post advertised silicone pads and guaranteed symptom improvement without scientific evidence (economic harm). Others encouraged their audience to forgo assessment by a doctor and pursue a pelvic floor physiotherapist instead (harmful inaction).

Regarding the reach of these 7 “probably harmful” posts, 5 of these posts came from Pinterest, with a modest reach of around 1000 to 4000 followers. The 1 Instagram post by a health blogger had the greatest reach with over 100 000 followers, and the 1 YouTube post by a doctor in Bangalore had the lowest reach with fewer than 200 subscribers and 400 views (Table 1).

## Discussion

In this novel investigation assessing the category and quantity of misinformation and resultant harm regarding POP on YouTube, Pinterest, and Instagram, our data show a significant difference in distribution of misinformation across platforms, and that misinformative posts regarding a benign condition can also result in harm, most frequently harmful inaction and economic harm and most frequently in Pinterest posts and least frequently in YouTube posts. These findings are congruent with a previous study by Morra et al. who found that YouTube is considered a reliable source of information on bladder pain syndrome, with more than 70% of videos receiving good or excellent global quality scores and more than 80% having no misinformation[6].

Limitations of the study are that the data were gathered from a small subset of posts and may not be generalizable to all medical-related content across

all social media platforms. Further limitations include the subjective nature of scoring and the selection of just 3 of dozens of social media platforms. While our methodology attempted to standardize review using multiple reviewers, subjectivity is naturally introduced when using nonbinary metrics for harm, where it is not possible to track which consumers of media pursued specific treatments and whether resultant harm ensued. Furthermore, while YouTube, Instagram, and Pinterest are among the top 15 social media platforms used globally, Facebook, the social media platform with the greatest number of daily and monthly users, was not included in our study because of the private nature of many pages and accounts that limited analysis of this magnitude.

## References

1. Ventola CL. Social media and health care professionals: benefits, risks, and best practices. *PT*. 2014;39(7):491–520. PMID: 25083128; PMCID: PMC4103576.
2. Johnson SB, Parsons M, Dorff T, Moran MS, Ward JH, Cohen SA, et al. Cancer misinformation and harmful information on Facebook and other social media: a brief report. *J Natl Cancer Inst*. 2022;114(7):1036–1039. doi: 10.1093/jnci/djab141. PMID: 34291289; PMCID: PMC9275772.
3. Barber MD, Maher C. Epidemiology and outcome assessment of pelvic organ prolapse. *Int Urogynecol J*. 2013;24(11):1783–1790. doi: 10.1007/s00192-013-2169-9. PMID: 24142054.
4. Pace LA, Herbert AS, Malik RD. Characteristics of pelvic organ prolapse content available on social media. *Neurourol Urodyn*. 2021;40(5):1165–1174. doi: 10.1002/nau.24673. PMID: 33834557.
5. Wardle C, Derakhshan H. Information disorder: toward an interdisciplinary framework for research and policy making. Council of Europe report DGI (2017)09. Available at: <https://rm.coe.int/information-disorder-toward-an-interdisciplinary-framework-for-research/168076277c>. Accessed January 29, 2023.
6. Morra S, Collà Ruvolo C, Napolitano L, La Rocca R, Celentano G, Califano G, et al. YouTube™ as a source of information on bladder pain syndrome: a contemporary analysis. *Neurourol Urodyn*. 2022;41(1):237–245. doi: 10.1002/nau.24802. PMID: 34559920.

## Conclusion

The rise of social media as a prominent source of healthcare information has outpaced research on its reliability and safety. Our research demonstrates the potential harms of various types of misinformation, even in a benign condition that impacts quality of life. Uniquely, our data help further categorize and quantify the specific type of misinformation across multiple social media platforms and sets the groundwork for analyzing the reach and impact of these posts. These data can help guide larger-scale policy changes, such as mandatory textual warnings underneath misinformative health posts, and help providers tailor medical education to best communicate patient diagnoses and treatment options.

**TABLE 1.**

Descriptive data for posts identified “Probably Harmful” medical claims regarding POP

In your opinion, are the medical claims within the social media post/video accurate?	Which section, if applicable, contained the misinformation?	What was misinformative about the post?	How would you categorize the medical claims made?
<b>Pinterest</b>			
Mostly false	Caption, Title, Hyperlink	Advertises silicone pad product (not evidence based)	Fabricated content
			Misleading content
Mostly false	Image, Hyperlink	Advertised book that proposes cure to urinary incontinence with Kegel exercises	Misleading content
			False context
Mostly false	Hyperlink	Claims that OB GYN doctors are not trained to diagnose pelvic floor dysfunction or prolapse and that pursuing OB GYN evaluation is not the best course of action	Fabricated content
			Misleading content
			Manipulated content
			False context
Mostly false	Image, Caption, Title, Hyperlink	Claims that starvation and dysbiosis are prominent risk factors for POP using out-of-context Nature article findings	Misleading content
			Manipulated content
			False connection
			False context
Mixture of both true and false	Hyperlinked website to post	Claims that tight hip flexors could cause pelvic discomfort	Fabricated content Manipulated content
<b>YouTube</b>			
Mixture of both true and false	Video	Claims that radiation and chemotherapy are nonsurgical options to treat POP	Fabricated content
<b>Instagram</b>			
Mixture of both true and false	Caption	Advertises electronic pelvic stimulator product (not evidence based)	Fabricated content
			Misleading content
			Manipulated content

POP: pelvic organ prolapse.

How would you categorize the type of harm caused by the medical claims?		Do you think there was purposeful intent by the creator to cause harm?	Number of likes or pins	Number of views	Publisher type	Speaker type	Number of followers or subscribers
Harmful inaction	Economic harm	Likely intentional	54		Commercial media/ industry	Industry	991
Harmful inaction							
Harmful inaction	Economic harm	Likely intentional			Commercial media/industry	Industry	1100
Harmful inaction							
Harmful inaction	Economic harm	Likely intentional			Consumer/patient	Health blogger	4100
Harmful inaction							
Harmful inaction	Economic harm	Uncertain if harm was intentional			Consumer/patient	Health blogger	4100
Harmful inaction							
Economic harm		Likely intentional			Consumer/patient	Physical therapist	2
Harmful action	Harmful interactions	Likely intentional	0	421	Doctor	Doctor	196
Harmful action							
Harmful action							
Harmful action	Economic harm	Likely intentional	839		Health/wellness group	Health blogger	111 000
Harmful action							
Harmful action							